



States has dropped by about 3 months in just the most recent 30 years, a move to a great extent ascribed to expanding body weights (Anderson et al., 2003).

Corpulence convolutes around half of all instances of polycystic ovarian disorder and most likely affects or compounds the natural ovulatory brokenness by enlarging both the insulin protection and the hyperandrogenemia show in this issue (Rittmaster et al., 1993). An affiliation free of polycystic ovarian disorder, be that as it may, has likewise been shown between truncal heftiness and unpredictable menses, proposing that corpulence itself may adversely affect fertility (Douchi et al., 2002).

Obesity and overweight are endemic worldwide. About 1 billion people are overweight of which 300 million are obese according to a WHO report, world widely (Warraich et al., 2009).

Obesity adversely influences reproductive function. It not only creates complications in conception and pregnancy but also increases the risk of miscarriage (Balen, et al., 2007; Pasquali, 2006).

## Materials and Methods

To investigate the relationship of infertility with weight and PCOs, 100 infertile female subjects were examined from May 2012 to June 2014. A comprehensive questionnaire was designed which contain almost all related information about infertility including background of reproductive physiology and chemical pathology. Questionnaire was filled in the presence of doctor or along with some personale. Levels of FSH, LH and prolactin hormone were also investigated along with PCOs test, in subjects. The data was divided into 3 groups i.e., Group I, II and III on the basis of age. Infertile women aged 19-42 years, having irregular or regular menstrual cycle were included, while women, who had conceived once or women had secondary infertility, aged less than 19 and more than 42 years were excluded.

## Statistical Analysis

Data examination was analyzed through the Statistical Package for the Social Sciences adaptation 11 (SPSS Inc., Chicago, IL, USA) for mean  $\pm$  standard error (SE). LSD test was applied, for finding differences among mean values. Statistical significance was assessed by Pearson's correlation test for Independence. The overall significance of differences between three groups (cases with PCOs, weight, systolic and diastolic blood pressure) was analyzed. A probability values (p value) less than 0.05 was considered significant. For categorical variables, the results were expressed as percentages.

## Results

GROUP I: In this group the mean age (yrs) was  $23.42 \pm 0.40$ , weight (Kg)  $50.02 \pm 0.96$ , while mean values of systolic and diastolic blood pressure (mm Hg) were  $108.75 \pm 2.27$  and  $73.75 \pm 1.88$  respectively. PCOs had been found in 25% subjects of this group (Table. 1).

GROUP II: The mean age (yrs) in this group was  $30.18 \pm 0.33$ , weight (Kg)  $66.29 \pm 0.62$ , systolic blood pressure (mm Hg)  $115.64 \pm 1.71$ , diastolic blood pressure (mm Hg)  $77.82 \pm 1.12$ . Subjects of this group had 45.45% of PCOs.

GROUP III: This group had the mean age (yrs)  $39.40 \pm 0.87$ , weight (Kg)  $86.76 \pm 1.43$ , systolic blood pressure (mm Hg)  $110.48 \pm 3.04$ , diastolic blood pressure (mm Hg)  $74.76 \pm 2.24$ . 57.14% of the subjects had found to have PCOs.

A significant correlation was observed between weight and PCOs ( $P = 0.01$ ) by statistical analysis of the data (Table. 2), while the relationship was found to be non-significant between weight and systolic ( $P = 0.23$ ) and weight and diastolic blood pressure ( $P = 0.17$ ).

## Discussion

Hippocrates 2500 years ago noted a pessimistic consequence of obesity on fertility, and in modern times obesity is associated with PCOS, chronic anovulation, infertility and menstrual irregularities (Wallach et al., 1987; Grodstein et al., 1994; Zaidi et al., 2009).

Many studies have been conducted in Africa, where the reported prevalence of infertility ranges from 9% in Gambia to 30 % in Nigeria (Gerrits, 2012). In China the prevalence of primary infertility found 9%, 10 – 15% in America, 16% in Siberia and 19% in Australia (Yi, Zeng, and Wu Deqing 2000). In various reports the rate of primary infertility has been reported 8 – 21.9 % (Safarinejad, 2008). By the WHO 8 – 12% infertility in the world is estimated (Dovom, et al., 2014).

Different factors including age, weight, PCOs, systolic and diastolic blood pressure that have been responsible for infertility in females were carefully studied in the present study. The prevalence and common causes of female infertility aged between 15 and 55 years had also been studied by Sule *et al.*, (2008).

**Table 1: Mean values of weight, Systolic blood pressure (mmHg), Diastolic blood pressure (mmHg) and PCOs in different age groups of specific female population of Karachi.**

Groups	Age of patient			Weight of patient			Systolic blood pressure (mmHg)	Diastolic blood pressure (mmHg)	Percent age of PCOS
	Max-Min	N	Mean ± SE	Max-Min	N	Mean ± SE	Mean ± SE	Mean ± SE	
I	25-19	24	23.42±0.403	58-38	24	50.021±0.969	108.75±2.277B	73.75±1.886A	25
II	35-26	71	30.18±0.330	78-60	55	66.291±0.624	115.64±1.717A	77.82±1.122A	45.45
III	42-27	5	39.40±0.872	100-80	21	86.762±1.432	110.48±3.045A B	74.76±2.249A	57.14

Means that do not share a letter in column are significantly different.

**Table 2: Coefficient of correlation (r) among various parameters of infertility.**

	Wt	Sys	Dia
Sys	R=0.121 ( <i>p</i> = 0.232)		
Dia	0.138 ( <i>p</i> = 0.170)	0.811 ( <i>p</i> = 0.000)	
PCO	-0.234 ( <i>p</i> = 0.019)	-0.008 ( <i>p</i> = 0.934)	0.037 ( <i>p</i> = 0.714)

Wt=Weight(Kg), Sys=Systolic blood pressure (mm/Hg),Dia=Diastolic blood pressure (mm/Hg), PCO=Polycystic Ovary.

In the present study minimum age limit were taken 19 years while the maximum age limit was 42 years. The findings of Ogunniyi et al. (1999) taken 15 years as lower limit of age because average female puberty starts at 13 years and around 15 years mostly pregnancy started and the upper limit of age was 47 years because menopause starts at the average age of 46 years. Demographical study noted that at the age of 35 in women birth rates starts to decline (Menken, et al., 1986). The average women at the age of 41 will deliver their last child ranging from 23-51 (Broekmans, et al., 2004).

Globally 1.3 billion people are estimated to be obese and overweight (World Health Organization, 2004). Obesity increased pregnancy complications and had a considerable impact on reproductive outcome (Pasquali, 2006; Balen et al., 2007). In current study 34% cases had body weight above 70 Kg. Conclusively mostly overweight subjects (n = 21) had PCOS, as ANOVA (Table 2) proved a significant relationship between weight and PCOs by statistical analysis of the data. The risk of infertility, conception rates, miscarriage rates, and pregnancy complications are increased in obese women with PCOs (Pettigrew and Hamilton-Fairley, 1997). Ovulatory cycles were regained in 12 out of 13 women and after losing 6.3 ± 4.8 Kg of weight, 11 subjects had conceived (Clark et al., 1995).

PCOs found to be most significant cause as in recent studies infertility incidence in the Fergana Valley was 16.8%, with polycystic ovaries' disease (PCOD) reported in Uzbekistan (Khaidarova, 2007). Current study also revealed that PCOs had profound effect in infertility, 43% of the cases had PCOs.

Infertility observed in patients having PCOs with higher rates approximately 75% (Hull, 1987; Stadtmauer and Oehninger, 2005). The current findings coincide with the results of Wallach, et al., (1988); Nestler, et al, (1989); Insler, et al., (1993) and Pasquali and Casimirri, (1993). They directly correlate hyper insulinaemia and hyper androgenaemia with obesity and PCOS.

## Conclusion

Current study has highlighted the link between obesity, infertility and adverse reproductive health outcome. Infertility is associated with several risk factors, multifactorial in nature. The age has profound effects on fertility along with weight in females. Obesity is increasingly prevalent health burdens upon modern society. All obese women are not infertile; however obesity and its negative impact upon fertility are well documented.

## References

- Anderson, S. E., Dallal, G. E., & Must, A. (2003). Relative weight and race influence average age at menarche: results from two nationally representative surveys of US girls studied 25 years apart. *Pediatrics*, *111*(4), 844-850.
- Bernasconi, D., Del Monte, P., Meozzi, M., Randazzo, M., Marugo, A., Badaracco, B., & Marugo, M. (1996). The impact of obesity on hormonal parameters in hirsute and nonhirsute women. *Metabolism-Clinical and Experimental*, *45*(1), 72-75.
- Balen, A. H., Anderson, R. A., & Policy & Practice Committee of the BFS. (2007). Impact of obesity on female reproductive health: British fertility society, policy and practice guidelines. *Human Fertility*, *10*(4), 195-206.
- Broekmans, F. J., Faddy, M. J., Scheffer, G., & te Velde, E. R. (2004). Antral follicle counts are related to age at natural fertility loss and age at menopause. *Menopause*, *11*(6 Part 1 of 2), 607-614.
- Cooper, T. G., Noonan, E., Von Eckardstein, S., Auger, J., Baker, H. W., Behre, H. M., & Vogelsong, K. M. (2010). World Health Organization reference values for human semen characteristics. *Human reproduction update*, *16*(3), 231-245.
- Clark, A. M., Ledger, W., Galletly, C., Tomlinson, L., Blaney, F., Wang, X., & Norman, R. J. (1995). Weight loss results in significant improvement in pregnancy and ovulation rates in anovulatory obese women. *Human Reproduction*, *10*(10), 2705-2712.
- Douchi, T., Kuwahata, R., Yamamoto, S., Oki, T., Yamasaki, H., & Nagata, Y. (2002). Relationship of upper body obesity to menstrual disorders. *Acta obstetrica et gynecologica Scandinavica*, *81*(2), 147-150.
- Dovom, M. R., Tehrani, F. R., Abedini, M., Amirshkari, G., Hashemi, S., & Noroozadeh, M. (2014). A population-based study on infertility and its influencing factors in four selected provinces in Iran (2008-2010). *Iranian journal of reproductive medicine*, *12*(8), 561.
- Frisch, R. E., & McArthur, J. W. (1974). Menstrual cycles: fatness as a determinant of minimum weight for height necessary for their maintenance or onset. *Science*, *185*(4155), 949-951.
- Gerrits, T. (2012). Biomedical infertility care in low resource countries: barriers and access. *Facts Views Vis Obgyn Monog*, *2*, 1-6.
- Grodstein, F., Goldman, M. B., & Cramer, D. W. (1994). Body mass index and ovulatory infertility. *Epidemiology*, 247-250.
- Holte, J., Bergh, T., Berne, C., & Lithell, H. (1994a). Serum lipoprotein lipid profile in women with the polycystic ovary syndrome: relation to anthropometric, endocrine and metabolic variables. *Clinical endocrinology*, *41*(4), 463-471.
- Holte, J., Bergh, T., Gennarelli, G., & Wide, L. (1994b). The independent effects of polycystic ovary syndrome and obesity on serum concentrations of gonadotrophins and sex steroids in premenopausal women. *Clinical endocrinology*, *41*(4), 473-481.
- Hull, M. G. R. (1987). Epidemiology of infertility and polycystic ovarian disease endocrinological and demographic studies. *Gynecological Endocrinology*, *1*(3), 235-245.
- Insler V, Shoham Z, Barash A, Koistinen R, Seppälä M, Hen M, Lunenfeld B, Zadik Z. (1993). Polycystic ovaries in non-obese and obese patients possible pathophysiological mechanism based on new interpretation of facts and findings. *Hum. Reprod*, *8*, 379-384.
- Imani, B., Eijkemans, M. J., Te Velde, E. R., Habbema, J. D. F., & Fauser, B. C. (1998). Predictors of patients remaining anovulatory during clomiphene citrate induction of ovulation in normogonadotropic oligoamenorrhic infertility. *The Journal of Clinical Endocrinology & Metabolism*, *83*(7), 2361-2365.
- Kirschner, M. A., Samojlik, E., Drejka, M., Szmaj, E., Schneider, G., & Ertel, N. (1990). Androgen-estrogen metabolism in women with upper body versus lower body obesity. *The Journal of Clinical Endocrinology & Metabolism*, *70*(2), 473-479.
- Khaidarova FA. (2007). Medico-biologic aspects of infertile marriage in the region of Fergana valley. *Problemi biologii i medicini*, 2007; 1: 119-122.
- Menken, J., Trussell, J., & Larsen, U. (1986). Age and infertility. *Science*, *233*(4771), 1389-1394.
- Nelson, L. R., & Bulun, S. E. (2001). Estrogen production and action. *Journal of the American Academy of Dermatology*, *45*(3), S116-S124.
- Nestler, J. E., Clore, J. N., & Blackard, W. G. (1989). The central role of obesity (hyperinsulinemia) in the pathogenesis of the polycystic ovary syndrome. *American Journal of Obstetrics & Gynecology*, *161*(5), 1095-1097.
- Norman, R. J., Davies, M. J., Lord, J., & Moran, L. J. (2002). The role of lifestyle modification in polycystic ovary syndrome. *Trends in Endocrinology & Metabolism*, *13*(6), 251-257.
- Noppa, H., & Bengtsson, C. (1980). Obesity in relation to socioeconomic status. A population study of women in Göteborg, Sweden. *Journal of Epidemiology & Community Health*, *34*(2), 139-142.

- Ogunniyi, S. O., Makinde, O. O., & Dare, F. O. (1990). Abortion-related deaths in Ile-Ife, Nigeria: a 12-year review. *African journal of medicine and medical sciences*, 19(4), 271-274.
- Pasquali, R. and Casimirri, F. (1993). The impact of obesity on hyperandrogenism and polycystic ovary syndrome in premenopausal women. *Clin. Endocrinol (Oxf)*, 39, 1-16.
- Pasquali, R. (2006). Obesity, fat distribution and infertility. *Maturitas*, 54(4), 363-371.
- Pedersen, S. B., Børghlum, J. D., Brixen, K., & Richelsen, B. (1995). Relationship between sex hormones, body composition and metabolic risk parameters in premenopausal women. *European journal of endocrinology*, 133(2), 200-206.
- Pettigrew, R., & Hamilton-Fairley, D. (1997). Obesity and female reproductive function. *British Medical Bulletin*, 53(2), 341-358.
- Pritts, E. A., Parker, W. H., & Olive, D. L. (2009). Fibroids and infertility: an updated systematic review of the evidence. *Fertility and sterility*, 91(4), 1215-1223.
- Rittmaster, R. S., Deshwal, N. A. T. A. S. H. A., & Lehman, L. (1993). The role of adrenal hyperandrogenism, insulin resistance, and obesity in the pathogenesis of polycystic ovarian syndrome. *The Journal of Clinical Endocrinology & Metabolism*, 76(5), 1295-1300.
- Safarinejad, M. R. (2008). Infertility among couples in a population-based study in Iran: prevalence and associated risk factors. *International journal of andrology*, 31(3), 303-314.
- Stadtmauer, L., & Oehninger, S. (2005). Management of infertility in women with polycystic ovary syndrome. *Treatments in endocrinology*, 4(5), 279-292.
- Sule, J. O., Erigbali, P., & Eruom, L. (2008). Prevalence of infertility in women in a southwestern Nigerian community. *African Journal of Biomedical Research*, 11(2).
- Wallach, E. E., Barbieri, R. L., Smith, S., & Ryan, K. J. (1988). The role of hyperinsulinemia in the pathogenesis of ovarian hyperandrogenism. *Fertility and sterility*, 50(2), 197-212.
- Wallach, E. E., Reid, R. L., & Van Vugt, D. A. (1987). Weight-related changes in reproductive function. *Fertility and Sterility*, 48(6), 905-913.
- Warraich, H. J., Javed, F., Faraz-ul-Haq, M., Khawaja, F. B., & Saleem, S. (2009). Prevalence of obesity in school-going children of Karachi. *Plos one*, 4(3), e4816.
- World Health Organization. (2004). Global strategy on diet, physical activity and health.
- Yi, Z., & Deqing, W. (2000). A regional analysis of divorce in China since 1980. *Demography*, 37(2), 215-219.
- Zaidi, S., Usmani, A., Shokh, I. S., & Alam, S. E. (2009). Ovarian reserve and BMI between fertile and subfertile women. *J Coll Physicians Surg Pak*, 19(1), 21-4.