

## CHEMICAL CONSTITUENTS FROM THE BIOACTIVE ETHYL ACETATE FRACTION OF *XANTHIUM STRUMARIUM* LINN.

<sup>1</sup>ANEELA WAHAB\*, <sup>1</sup>AMINA SULTANA, <sup>2</sup>KHALID M. KHAN, <sup>2</sup>SHAZIA SHAH, <sup>3</sup>SIKANDAR KHAN SHERWANI AND <sup>1</sup>SYED SAJJAD HAIDER

<sup>1</sup>Department of Chemistry, Federal Urdu University of Arts, Science and Technology, Gulshan-e-Iqbal, Karachi, Pakistan

<sup>2</sup>H.E.J Research Institute of Chemistry, International Center for Chemical and Biological Sciences, University of Karachi, Pakistan

<sup>3</sup>Department of Microbiology, Federal Urdu University of Arts, Science and Technology, Gulshan-e-Iqbal, Karachi, Pakistan

\*Corresponding author: e-mail: [anilawahab@fuuast.edu.pk](mailto:anilawahab@fuuast.edu.pk)

### خلاصہ

زینتھیم اسٹرومیم کی ایٹھائیل ایسی ٹیسٹ موجودگی مخالف مائیکرو حیاتیاتی اور آکسی ڈیٹھ خصوصیات کی حامل پائی گئی۔  $EC_{50} = 937 \text{ ug/ml}$  اہیت کی حامل مخالف Oxident عمل کو ظاہر کرتی ہے۔ اس کاربک تیزاب یا معیار کی مخلول کے برخلاف۔ یہ درمیانی درجہ سے اعلیٰ درجہ کا میکرو حیاتیاتی مخالف رو یہ اختیار کرتی ہے جو 12-34 mm کا مخالف علاقہ بناتی ہے۔ Pentacyclic terpenes جس میں B-armyryn اور Oleanolic تیزاب اور Aromatic تیزاب جسمیں Ferulic تیزاب شامل ہے۔ اس حیاتیاتی اہیت کے اتھائیل لیسٹیٹ جو کہ Strumarium Xanthinm میں سے حاصل کیئے جاتے ہیں۔ اس مرکب کی شناخت کو مختلف اسپیکٹرو اسکوپک طریقہ کار سے جو کہ IR – EIRS اور H-NMR طریقہ کار سے حاصل کیا گیا ہے۔

### Abstract

The ethyl acetate fraction of *Xanthium strumarium* Linn. was found possessing antioxidant and antimicrobial potential. It showed significant *in vitro* antioxidant activity ( $EC_{50} = 937 \text{ } \mu\text{g/mL}$ ) in contrast of ascorbic acid as standard. It also exhibited moderate to high antibacterial and antifungal activities having zone of inhibition in range of 12-34 mm against pathogenic microorganism. Two pentacyclic triterpenes  $\beta$ - amyryn (1), oleanolic acid (2) and an aromatic acid, ferulic acid (3) have been isolated from this bioactive ethyl acetate (EtOAc) fraction of *Xanthium strumarium* Linn. The structures of these compounds were elucidated through various spectroscopic techniques such as IR, EIMS and <sup>1</sup>H-NMR techniques.

### Introduction

*Xanthium strumarium* L. belongs to family compositeae (Winter *et al.*, 1969). The genus *Xanthium* consist of 25 species and distributed near temperate and tropical region of world (Han *et al.*, 2007). The members of genus *Xanthium* L. used to cure herpes, fever, nasal sinusitis, urticaria, headache, arthritis, cancer, and scrofula (Ravindra, 2003). *X. strumarium* L. has been used as a most important plant in herbal medicine in Malaysia, India and China (Prajapati and Purohit, 2003). Previously sesquiterpenoids, alkaloids, and sesquiterpene lactones (Gupta and Gupta, 1975) and Carbohydrates (Craig *et al.*, 1976) have been reported from this species. Chemical constituents of this plant possess antimicrobial activity (Jawad *et al.*, 1988), antifungal activity (Sastri, 1976) and cytotoxic activity (Kim *et al.* 2003). Fruits are used to treat small-pox; and contain anti-inflammatory agent (Sastri, 1976) and anti-hyperglycemic agent (Hsu *et al.* 1999). Seeds have toxic effects due to choline and hydroquinone (Sastri, 1976). Biological activities of *X. strumarium* extracts were also reported such as anticancer, antitrypanosomal (Anooj and Ajay, 2010), antimalarial (Joshi *et al.*, 1997), hepatoprotective (Panday and Rather, 2012) and analgesic properties (Han *et al.*, 2007).

The ethnopharmacological and chemotaxonomic importance of the genus *Xanthium* stimulated us to further investigate on *X. strumarium*. Previously we have reported antimicrobial, antioxidant potential and chemical constituents of n-hexane, dichloromethane and n-butanol fraction (Aneela *et al.*, 2012). Herein, we have investigated ethylacetate fraction of *X. strumarium* to determine its anti-bacterial, anti-fungal and anti-oxidant activities and possibly active substances.

## Experimental

### Materials and Methods

**General:** Gallenkamp melting point apparatus was used to determine melting points. Silica gel (70-230 mesh, Merck) was used for column chromatography (CC), whereas, Pre-coated silica gel (GF-254) was used for TLC. IR spectra were scanned on Jasco-302-A spectrophotometer. EIMS spectra were recorded on Mass spectrometer (Jeol-JMS HX-110). Bruker spectrometer (500 and 300 MHz) was used to record <sup>1</sup>H-NMR spectra. The chemical shift values were reported in  $\delta$  (ppm) relative to tetramethylsilane (TMS) as an internal standard. The coupling constant ( $J$ ) values were given in Hz.

**Plant Material:** The aerial part of plant material *X. strumarium* were collected from Lakimarwat (NWFP) Pakistan and identified by Dr. Sahar, Department of Botany, University of Karachi (voucher specimen G.H. No.86398, No. 01).

**Extraction and Isolation:** The freshly collected shade dried aerial parts of plant material *X. strumarium* (10 kg) were ground and extracted with methanol (15 L). The resulting extract was concentrated under vacuum to obtain a crude extract (250 g) which was then successively extracted with *n*-hexane (13 g), dichloromethane (14 g), ethyl acetate (97 g) and *n*-butanol (21 g). The ethyl acetate fraction (97 g) was subjected to CC over silica gel and eluted successively with *n*-Hexane and *n*-Hexane-ethylacetate in an increasing order of polarity to obtained 16 sub-fractions (A-1 to A-16). The sub-fraction A-3, eluted with solvent *n*-hexane-ethylacetate (9:1) was a semi-pure compound which was further purified by preparative TLC (Hexane: ethylacetate, 9:1) to give compound **1** ( $\beta$ -amyryn). The sub-fraction A-6 was subjected to CC (with an increasing order of polarity, *n*-hexane, *n*-hexane-EtOAc) which gave ten sub-fractions (A-6-1 to A-6-10). The sub-fraction A-6-3 obtained from solvent (*n*-hexane-ethylacetate, 8:2) was kept overnight in a mixture of solvent (MeOH:CHCl<sub>3</sub>, 1:1) to obtain compound **2** (oleanolic acid). The sub-fraction A-9 was subjected to CC (with an increasing order of polarity of solvent CHCl<sub>3</sub>, CHCl<sub>3</sub>-MeOH) gave six sub-fractions (A-9-1 to A-9-6). Fraction A-9-3 was obtained from solvent system (CHCl<sub>3</sub>-MeOH, 9:1) to afford compound **3** (ferulic acid).

**Characterization of  $\beta$ -Amyryn (1):** Colourless solid (6.0 mg). M.P., 195- 196° C; IR (KBr)  $\nu_{\max}$  cm<sup>-1</sup>: 3352 (OH), 1648 (C=C); EI-MS  $m/z$  (rel. int.,%): 426 [M]<sup>+</sup> (49), 411 (15), 408 (10), 393 (10), 218 (100), 207 (42), 203 (53), 189 (60), 175 (24), 147 (25), 135 (45), 119 (27), 109 (41) and 69 (55); HR-EIMS  $m/z$ : 426.0100 (calcd. for C<sub>30</sub>H<sub>50</sub>O, 426.3862); 411.3412 [C<sub>29</sub>H<sub>47</sub>O]<sup>+</sup>, 408.1502 [C<sub>30</sub>H<sub>48</sub>]<sup>+</sup>, 393.223 [C<sub>29</sub>H<sub>45</sub>]<sup>+</sup>, 218.0420 [C<sub>16</sub>H<sub>26</sub>]<sup>+</sup>, 207.3301 [C<sub>14</sub>H<sub>23</sub>O]<sup>+</sup>, 203.1354 [C<sub>15</sub>H<sub>23</sub>]<sup>+</sup>, 189.3010 [C<sub>14</sub>H<sub>21</sub>]<sup>+</sup>, 175.0132 [C<sub>13</sub>H<sub>19</sub>]<sup>+</sup>, 147.1945 [C<sub>11</sub>H<sub>15</sub>]<sup>+</sup>, 135.1093 [C<sub>10</sub>H<sub>15</sub>]<sup>+</sup>, 119.2203 [C<sub>9</sub>H<sub>11</sub>]<sup>+</sup>, 109.0110 [C<sub>8</sub>H<sub>13</sub>]<sup>+</sup> and 69.2015 [C<sub>3</sub>H<sub>9</sub>]<sup>+</sup>; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  5.15 (1H, t,  $J$  = 4.5 Hz, H-12), 3.09 (1H, dd,  $J$  = 11.0, 4.0 Hz, H-3 $\alpha$ ), 1.08 (3H, s, CH<sub>3</sub>), 1.00 (3H, s, CH<sub>3</sub>), 0.99 (3H, s, CH<sub>3</sub>), 0.94 (3H, s, CH<sub>3</sub>), 0.87 (3H, s, CH<sub>3</sub>), 0.86 (3H, s, CH<sub>3</sub>), 0.85 (3H, s, CH<sub>3</sub>), and 0.78 (3H, s, CH<sub>3</sub>). All the spectral data were in agreement with values reported in literature (Malik *et al.*, 2005).

**Characterization of Oleanolic acid (2):** Colourless needles (4.0 mg); MP: 195-197°C; IR (KBr)  $\nu_{\max}$  cm<sup>-1</sup>: 3410 (OH), 2789 (COOH), 2768 (C-H), 1710 (C=O), 1625 (C=C), and 1066 (C-O); EI-MS  $m/z$  (rel. int.): 456 [M]<sup>+</sup> (4), 248 (100), 207 (23), 203 (31), 189 (10) and 133 (21); HR-EIMS  $m/z$ : 456.3410 (calcd for C<sub>30</sub>H<sub>48</sub>O<sub>3</sub>, 456.3530), 248.1730 [C<sub>16</sub>H<sub>24</sub>O<sub>2</sub>]<sup>+</sup>, 207.1643 [C<sub>14</sub>H<sub>23</sub>O]<sup>+</sup>, 203.2735 [C<sub>15</sub>H<sub>23</sub>]<sup>+</sup>, 189.1549 [C<sub>14</sub>H<sub>21</sub>]<sup>+</sup> and 133.1081 [C<sub>10</sub>H<sub>13</sub>]<sup>+</sup>; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz):  $\delta$  5.30 (t,  $J$  = 3.6 Hz, H-12), 3.42 (1H, dd,  $J$  = 14.0, 4.0 Hz, H-3 $\alpha$ ), 2.80 (1H, dd,  $J$  = 14.1, 4.6 Hz, H-18), 1.20 (3H, s, CH<sub>3</sub>), 1.15 (3H, s, CH<sub>3</sub>), 0.97 (3H, s, CH<sub>3</sub>), 0.94 (3H, s, CH<sub>3</sub>), 0.86 (3H, s, CH<sub>3</sub>), 0.85 (3H, s, CH<sub>3</sub>), and 0.75 (3H, s, CH<sub>3</sub>). The studied spectral data coincided with those in literature (Misra *et al.*, 1981).

**Characterization of Ferulic acid (3):** Colourless solid (3.0 mg); M.P., 168-172 °C; IR (KBr)  $\nu_{\max}$  cm<sup>-1</sup>: 3345 – 2578 (OH, COOH), 1675 (C=O), 1619 and (C=C); EI-MS  $m/z$  (rel. int. %): 194 [M]<sup>+</sup> (10), 149 (24), 163 (32), 123 (19) and 92 (5); HR-EIMS  $m/z$ : 194.0136 (calcd. for C<sub>10</sub>H<sub>10</sub>O<sub>4</sub>, 194.2110), 149 [C<sub>9</sub>H<sub>9</sub>O<sub>2</sub>]<sup>+</sup>, 163 [C<sub>9</sub>H<sub>7</sub>O<sub>3</sub>]<sup>+</sup>, 123 [C<sub>7</sub>H<sub>7</sub>O<sub>2</sub>]<sup>+</sup> and 92 [C<sub>6</sub>H<sub>4</sub>O]<sup>+</sup>; <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 500 MHz):  $\delta$  7.48 (1H, d,  $J$  = 14.5 Hz, H-7), 7.21 (1H, d,  $J$  = 7.8 Hz, H-6), 7.15 (1H, br. s, H-2), 6.59 (1H, d,  $J$  = 7.9 Hz, H-5), 6.01 (1H, d,  $J$  = 14.5 Hz, H-8) and 3.89 (3H, s, OCH<sub>3</sub>). The given spectral data agreed the literature values (Li *et al.*, 1999).

### Biological Assay

**Preparation of media:** The media used for culturing of bacterial strain was Muller Hinton agar and Muller Hinton broth s and fungal strains were culturing on Sabour dextrose agar (SDA) (Smyth *et al.*, 2011).

**Screening of Antibacterial activity:** Antibacterial activity of ethylacetate fraction was carried out by agar well diffusion method. Muller Hinton broth was autoclaved for 2 h and then culture (10  $\mu$ l) was transferred into the wells (Perez *et al.*, 2009). Test fraction (10 mg/mL) was incubated at 28 °C for 48 hr and then measured zone of inhibition. The standard drug gentamicin was used.

**Screening of Antifungal activity:** Ethyl acetate fraction was screened for antifungal activity by agar-well diffusion method. The media was shifted aseptically into each SDA plates (Wuthi *et al.*, 2011). Test fraction (10 mg/mL) were incubated for 48 hr. at 28  $\pm$  2°C and then measured zone of inhibition. The standard drug gresiofulvin was used.

**Minimum Inhibitory Concentration (MIC):** MIC was calculated by broth dilution method. Stock solution (100 mg/mL) of extract was made in dist. H<sub>2</sub>O. Further two fold serial dilution of extract was made in 100  $\mu$ L broth afterward 10  $\mu$ l culture matched with 0.5 Mac Farland index was added to each well. One well as culture control while other as antibiotic control. All plates were incubated at 37 °C for 24 hours. The MIC was establish when no visible growth was observed (Sherwani *et al.*, 2013).

**Screening of anti-oxidant activity:** Ethylacetate fraction of *X. strumarium* was screened for antioxidant potential by the method of Lee *et al.* (1998). DPPH (1,1-diphenyl-2-picrylhydrazile) solution was made in ethanol. 10  $\mu$ L EtOAc fraction and 90  $\mu$ L DPPH were added in 96-well microtiter plates and incubated for 30 min. at 37 °C. Absorbance was measured through spectrophotometer at wavelength of 515 nm. % inhibition of radicals was determined by following formula.

$$\% \text{ Inhibition} = \frac{(\text{absorbance of the control} - \text{absorbance of the test sample})}{\text{Absorbance of the control}} \times 100$$

## Results and Discussion

Compound **1** ( $\beta$ -amyrin) was isolated as colorless solid. The molecular ion peak at  $m/z = 426$  was observed by EI-MS and molecular formula C<sub>30</sub>H<sub>50</sub>O was obtained by HR-EIMS. Two peaks at  $m/z$  218 and 207 was obtained in EIMS due to Diels Alder fragmentations (Misra *et al.*, 1981). The absorption band in IR spectrum at 3352 cm<sup>-1</sup> is due to -OH group and absorption band at 1648 cm<sup>-1</sup> supported olefinic carbons (C=C). <sup>1</sup>H-NMR displayed eight singlet of methyl proton at  $\delta$  1.08 -0.78 and a triplet at  $\delta$  5.15 ( $J = 4.5$  Hz) is of olefinic protons. A double doublet (dd) at  $\delta$  3.09 ( $J = 11.0$  and 4.0 Hz) is the characteristic feature of H-3 $\alpha$  (Amzad and Zhari, 2013). All these data showed that compound **1** belongs to pentacyclic triterpene of  $\Delta^{12}$   $\beta$ -amyrin series (Malik *et al.*, 2005).

Compound **2** (oleanolic acid) was isolated as colorless needles. The [M]<sup>+</sup> peak at  $m/z = 456$  was observed by EI-MS and molecular formula C<sub>30</sub>H<sub>48</sub>O<sub>3</sub> was observed by HR-EIMS. Mass fragments at  $m/z$  248 and 207 due to retro Diels Alder fragmentation further supported that compound belongs to the pentacyclic triterpenes (Misra *et al.*, 1981). The IR spectrum showed the absorption band at 2789 cm<sup>-1</sup> due to presence of -COOH and absorption bands at 3410 due to presence of -OH group. while the presence of -COOH group at C-17 was indicated by the mass fragments at  $m/z$  203, 189 and 133 whereas -OH group at C-3 with  $\beta$  -orientation is on biogenetic ground. <sup>1</sup>H-NMR spectrum displayed seven methyls singlet at  $\delta$  1.20-0.75 and a triplet at  $\delta$  5.30 ( $J = 3.6$  Hz) of olefinic proton and a dd at  $\delta$  3.42 ( $J = 14.0$  Hz, 4.0 Hz) of H-3 $\alpha$  oxymethine (Amzad and Zhari, 2013). It was analyzed that compound **2** also belongs to series of  $\Delta^{12}$   $\beta$ -amyrin (Misra *et al.*, 1981).


Compound **3** (ferulic acid) was isolated as colorless solid. EIMS displaying [M]<sup>+</sup> peak at  $m/z = 194$ . Mass fragment at  $m/z$  149 was observed due to loss of -COOH group, at  $m/z$  163 due to loss of -OCH<sub>3</sub> and at  $m/z$  123 due to loss of olefinic linkage. Absorption band of IR at 1619, 3345 and 2578 cm<sup>-1</sup> exhibited of -C=C, -OH and -COOH group. <sup>1</sup>H-NMR displayed a singlet of methoxy group at  $\delta$  3.89 and two doublet of *trans* olefinic protons at  $\delta$  7.48 and 6.01 ( $J = 14.5$  Hz). A downfield singlet at  $\delta$  7.15 (s, H-2) and two doublet at 7.21 and 6.59 ( $J = 7.9$  Hz) showed *meta* substituted -OCH<sub>3</sub> and *para* substituted -OH in aromatic ring (Muhammad *et al.*, 2011 and Li *et al.*, 1999).

The ethylacetate fraction was evaluated for antimicrobial and antioxidant potential. Antibacterial activity was screened against *B. cereus*, *B. subtilis*, *B. thuringiensis*, *C. coli*, *E. aerogenes*, *K. pneumonia*, *S. typhi*, *S. paratyphi A*, *S. paratyphi B*, and *Vibrio cholera* and antifungal activity was evaluated against *A. flavus*, *A. niger*, *Helminthosporium*, *M. canis*, *Penicillium sp.*, and *Penicillium sp.* (Table 1). The ethylacetate fraction showed excellent antibacterial activity within the range of 17 to 34 mm and MIC value ranges 52 to 200 mg/mL. Ethylacetate fraction also showed significant antifungal activity (12-19 mm) and MIC value was observed within the range of 100-320 mg/mL.


The ethylacetate fraction also showed significant antioxidant potential (Table 2). DPPH free radical showed 71 % inhibition and the result of EC<sub>50</sub> was 938  $\pm$  0.5  $\mu$ g/ml.

**Conclusion**


The present study concludes that ethylacetate fraction of *Xanthium strumarium* is biologically active and has potential to inhibit the growth of various pathogenic Gram positive, Gram negative bacteria and is also minimizing fungal infections. It is also exhibited good antioxidant potential.



$\beta$ -Amyrin (1)



Oleanolic acid (2)



Ferulic acid (3)

**Table.1** *In vitro* anti-bacterial and anti-fungal activity of ethyl acetate fraction of *X. strumarium* Linn.

Organism	Zone of inhibition (mm)	Standard drugs	MIC (mg/mL)
<b>Gram positive bacteria</b>			
		<b>Gentamicin</b>	
<i>Bacillus cereus</i>	26 ± 2	>15	88
<i>Bacillus subtilis</i>	34 ± 0	>15	120
<i>Bacillus thuringiensis</i>	30 ± 0	>15	102
<b>Gram negative bacteria</b>			
<i>Enterobacter aerogenes</i>	17 ± 1	>15	200
<i>Klebsiella pneumoniae</i>	22 ± 1	>15	56
<i>Salmonella typhi</i>	18 ± 1	>15	89
<i>Salmonella paratyphi A</i>	19 ± 1	>15	90
<i>Salmonella paratyphi B</i>	20 ± 0	>15	52
<i>Campylobacter coli</i>	19 ± 0	>15	200
<i>Vibrio cholerae</i>	19 ± 1	>15	160
<b>Fungi</b>			
		<b>Gresiofulvin</b>	
<i>Microsporium canis</i>	12 ± 2	>12	312
<i>Aspergillus flavus</i>	19 ± 1	>12	320
<i>Aspergillus niger</i>	15 ± 0	>12	100
<i>Penicillium sp</i>	12 ± 1	>12	149
<i>Rhizopus</i>	16 ± 1	>12	300
<i>Helminthosporum</i>	14 ± 0	>12	212

**Table-2:** *In vitro* anti-oxidant activity of Ethyl acetate fraction of *X. strumarium* Linn.

Sample	% inhibition	EC <sub>50</sub> (ug/ml)	Standard drug
Ethyl acetate fraction	71 %	938 ± 0.5	Ascorbic acid

**References**

- Amzad, H. and Zhari, I. (2013). Isolation and characterization of triterpenes from the leaves of *Orthosiphon stamineus*, *Arabian Journal of Chemistry* 6(3): 295-298.
- Aneela, W., Khalid, M. K., Amina, S. and Aisha, I. (2012). Chemical investigation of *Xanthium strumarium* Linn. and Biological Activity of its Different Fractions, *Journal of pharmaceutical Research* 4: 1984-1987
- Anooj K. and Ajay, K. S. (2010). Phytopharmacological Review of *Xanthium strumarium* L. (Cocklebur). *International Journal of Green Pharmacy* 4:129-139.
- Craig, J.C., Mole, M. L. Billets, S. and Farouk, E.F. (1976). Isolation and Identification of the Hypoglycemic Agent, Carboxyatractylate from *Xanthium strumarium*. *Phytochemistry* 15: 1178.
- Gupta, K.C. and Gupta, D.R. (1975). Isolation of an Antibacterial compound, Xanthumin from the Leaves of *Xanthium strumarium* Linn, *Journal of Indian Chemical Society* 52:1224.-1225.
- Han, T., Li, H.L., Zhang, Q. Y. and Qin, L. P. (2007). Bioactivity-Guided Fraction for Anti-inflammatory and Analgesic Properties and Constituents of *Xanthium strumarium* Linn. *Phytomedicine* 14:825
- Hsu, F. L., Chen, Y. C. and Cheng, J. T. (1999). Caffeic Acid as Active Principle from the Fruit of *Xanthium strumarium* to Lower Plasma Glucose in Diabetic Rats. *Planta Medica* 66: 228-230.
- Jawad, A. I. M., Mahmoud M. J. and Al-Naib, A. (1988). Antimicrobial Activity of *Xanthium strumarium* Extracts. *Fitoterapia* LIX (3): 220-221.
- Joshi, S. P., Rojatar, S. R. and Nagasampagi, B. A. (1997). Antimalarial Activity of *Xanthium strumarium*. *Journal of Medicinal and Aromatic Plant Sciences* 19: 366-368.
- Kim, Y. S. Kim, J. S. and Kim, Y.K. (2003). Two Cytotoxic Sesquiterpene Lactones from the Leaves of *Xanthium strumarium* and their In vitro Inhibitory Activity on Farnesyltransferase. *Letter Planta Medica* 69:375.-377.
- Lee, S. K., Zakaria, H., Chung, H., Luyengi, L. and Pezzuto, J. M. (1998). Evaluation of the Antioxidant Potential of Natural Products. *Combinatorial Chemistry and High Throughput screening* 1: 35-46.
- Li, Y. M., Jiang, S. H. and Zhu, D. Y. (1999). Iridoid Glycosides from *Scrophularia ningpoensis*. *Phytochemistry* 50:101-104.
- Malik, A., Ferheen, S. and Ahmed, E. (2005). Phytochemical Studies on *Haloxylon salicornicum*. *Journal of Chemical Society of Pakistan* 27(2): 219-222.
- Misra, T. N., Singh R. S. and Ojha, T. N. (1981). Chemical Constituents of *Hyptis suaveolens*. Part 1. Spectral and Biological Studies on a Triterpene Acid. *Journal of Natural Product* 44: 735-738.
- Muhammad, S. A., Muhammad, I. A., Zeeshan, A. and Alain, F. K. W. (2011). Vogelate: A new long-chained Ester from *Erythrina vogelii* Hook (Fabaceae) of Cameron. *Journal of Chemical Society of Pakistan* 33(3): 412-416.
- Pandey, D. P. and Rather, M. A. (2012). Isolation and Identification of Phytochemicals from *Xanthium strumarium* L., *International Journal of Chem Tech Research* 4(1):266-271.
- Perez, C., Paul, M., and Bazerque, P. (2009). An Antibiotic Assay by the Agar Well Diffusion Method. *Acta Biologie Et Medicine Experimenta* 15: 113-115.
- Prajapati, N. D. and Purohit, S. S. (2003). *A Hand Book of Medicinal Plants*, A complete Source Book, Agribios (India), pp. 548.
- Ravindra, S., (2003). *Medicinal Plants of India*. An Encyclopaedia. Daya Publishing House, Delhi (India), pp.263.
- Sastri, B. N. (1976). *The Wealth of India*, CSIR, New Delhi (India). Vol. 10, pp.1.
- Sherwani, S. K., Bokhari, T. Z., Nazim, K. and Urooj, S. K. (2013). Qualitative Phytochemical Screening and Antifungal Activity of Lanicapapaya Leaf Extract Against Human and Plant Pathogenic Fungi. *International Research Journal of Pharmacy* 4: 83-86.
- Smyth, R. W., Bengtsson, S., Cloke, J. and Kahlmeter, G. (2011). Performance evaluation of M.I.C. Evaluator™strip(Oxid) and comparison with the BSAC reference methods. <http://www.oxidhai.com/wp-content/uploads/2011/04/ECCMID-2008-Swedish-poster.pdf>
- Winter, T. E., Geissmann, T.A. and Safir, D. (1969). Sesquiterpene Lactones of *Xanthium* Species. Xanthanol and Isoxanthanol and Correlation of Xanthinin with Ivalbin. *The Journal of Organic Chemistry* 34:153.
- Wuthi, U., Mansuang, and Omboon, V. (2011). In vitro Effectiveness of *Acacia concinna* Extract against Dermatocytoc Pathogens. *Pharmacognosy Journal* 3: 69-74.